Mediterranean Journal of Pharmacy & Pharmaceutical Sciences
ISSN: 2789-1895 online
www.medjpps.com
doi: 10.5281/zenodo.21852547
ISSN: 2958-3101 print

SHORT COMMUNICATION article

# Chemical composition and amino acid analysis of the seeds of the milk thistle plant (*Silybum marianum Gaertn* L.) widespread in Tulkarem, Palestine

**Khaled Taha Muhammad Abu Thaher**

College of Law and Forensic Sciences - Department of Forensic Sciences, Al-Istiqlal University (Palestinian Academy for Security Sciences), Jericho,Palestine

**Received:** 11-05-2026, **Accepted**: 09-08-2026, **Published online:** 10-08-2026

Copyright© 2026. This open-access article is distributed under the *Creative Commons Attribution License*, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**HOW TO CITE THIS**

Abu Thaher KTM. Chemical composition and amino acid analysis of the seeds of the milk thistle plant (*Silybum marianum Gaertn* L.) widespread in Tulkarem, Palestine. Mediterr J Pharm Pharm Sci. 2026; 6(3): 65-71. [Article number: 262]. https://doi.org/10.5281/zenodo.21852547

***Keywords:*** Amino acids, flavonoids, phenols, phytochemical investigation, Flora of Palestine, tannins

**Abstract:** In this study, the milk thistle (*Silybum marianum* Gaertn L.), widespread in Palestine, was selected to study the basic qualitative chemical composition of aqueous, ethanol, chloroform, and hexane extracts from its seeds, and to identify the active compounds present. The results of the phytochemical analysis showed that the seeds of the plant (*Silybum marianum* Gaertn L.) contain a full range of chemically active compounds in all its extracts, such as: flavonoids, amino acids, carbohydrates, sterols, saponins, anthraquinones, phenolic compounds, steroids, terpenoids, tannins, alkaloids, anthocyanins, quinones, and glycosides. For the first time in Palestine, seventeen amino acids were identified in *Silybum marianum* Gaertn L. plant seeds using phytochemical analysis, paper and thin-layer chromatography, and standard amino acid assays. As a result of the phytochemical analysis, a range of amino acids were identified in the plant seeds, including: arginine, leucine, valine, lysine, phenylalanine, isoleucine, threonine, histidine, glutamic acid, glycine, serine, proline, aspartic acid, alanine, tyrosine, methionine, cysteine.

## Introduction

The use of medicinal plants dates back to the dawn of human history. In recent years, the use of medicinal plants containing phenolic compounds has increased due to their diverse medicinal effects, low cost, and lack of side effects compared to manufactured drugs [1, 2]. Especially for plants that are globally important in the fields of nutrition and the treatment of various diseases, the determination of the qualitative chemical composition of the active substances in their parts, especially local species that have not been thoroughly studied in the country, has attracted the attention of researchers and scientists [2].

*Silybum marianum* (L.) Gaertn, commonly known as Milk Thistle, belongs to the Asteraceae family. It is an annual or biennial herbaceous plant native to the Mediterranean basin and now widespread throughout the world [3]. Its seeds contain a range of active compounds, and the plant is used to treat liver problems and other ailments. The leaves are spiny and have lateral spots around the veins. Milk thistle is self-pollinating and has pink or purple flowers, blooming in May and June. It has been used in traditional folk medicine for over 2000 years, most commonly in the treatment of liver diseases such as liver toxicity, cirrhosis, and hepatitis [4]. **Figure 1**. Milk thistle is cultivated commercially as a medicinal plant in Europe, Egypt, China, and Argentina [5]. On the other hand, based on the results of studies and research, milk thistle is currently considered an interesting crop for bioenergy production in the Mediterranean environment [6] and a source for biodiesel production [7].

ISSN: 2789-1895 online
www.medjpps.com
doi: 10.5281/zenodo.21852547
ISSN: 2958-3101 print



**Figure 1:** Plants, flowers and seeds of *Silybum marianum* L.

Extracts obtained from milk thistle are used for multiple medicinal purposes, such as anti-diabetic, liver and heart protection, cholesterol reduction, blood pressure reduction, anti-hemorrhagic, anti-inflammatory and antiviral, antispasmodic, nerve protection, anti-cancer, antioxidant, and having a protective effect against Alzheimer's [3, 8-12]. Milk thistle has therapeutic effects, which are closely related to the presence of a flavonoid compound called silymarin [13]. This compound stimulates liver regeneration and is recognized as an antioxidant and anti-inflammatory agent, and is therefore effective in treating food poisoning, hepatitis, and liver cirrhosis [14]. Recently, the hepatoprotective activity of silymarin, which is based on its antioxidant properties, has been documented, and it has become used in the manufacture of plant-based pharmaceuticals such as (Lagalo, Flavobion) used in the treatment of mycotoxicosis that affects the liver (Amanita sp.) [15]. Silymarin has been found in the highest concentration in the fruit of the plant, but it is also present in the leaves and seeds. At the beginning of 2014, silymarin extracted from milk thistle seeds was ranked sixth among the top 20 best-selling dietary supplements in the health and natural food market, and ranked twelfth among the top 40 herbal supplements in the plant-based products market in the United States [16, 17].

In different countries, the leaves, stems, flowers, and seeds of Milk thistle have been used for various purposes. Milk thistle seeds contain silymarin, bioflavonoids, flavonoids, amino acids, glutamic acid, fumaric acid, saponins, sterols, tocopherols, oils, starches, gum, minerals, tannins, fatty acids and polyphenolic components [18], and the oil extracted from these seeds can be used as a treatment for many diseases, including viral hepatitis and cirrhosis of the liver [19, 20]. Studies have shown that seeds contain many active substances beneficial to health, such as protein, with a valuable content of essential amino acids, and some active phytochemicals that possess pharmacological properties such as antioxidants or anti-cancer and anti-free radical effects [21]. In Palestine, only a few studies have been conducted on the uses of milk thistle in traditional medicine [22, 23]. Furthermore, adding milk thistle seed flour to food products and dietary supplements improves their absorption. Experts recommend consuming these products daily to help prevent diseases such as cardiovascular disease and some types of cancer. Therefore, studying the phytochemicals of the active ingredients in seeds is a crucial strategy [21]. The current study aimed to study the qualitative phytochemical composition of (*Silybum marianum* L.) seeds and to determine the qualitative chemical composition of the seeds in terms of amino acids. In addition, to the best of the researcher's knowledge, the qualitative phytochemical composition of the seeds of the wild *Silybum marianum* plant in Palestine has not been explored.

## Materials and methods

*Plant collection:* Samples of milk thistle (*Silybum Marianum* L.) seeds were carefully collected at maturity from the area of Far'un village (32°17′10″N 35°01′23″E) in the city of Tulkarm in the West Bank of Palestine during October 2025. The seeds were then left to dry naturally at room temperature for 30 days. After that, the seeds were ground into a fine powder. We will use this powder for qualitative chemical analysis of the active ingredients and to detect amino acids. The plant was formally identified as (*Silybum Marianum* L.) by Dr. Rami Al-Qaisi from the National Agricultural Research Center - MOA.

*Preparation of (Silybum marianum L.) seed extracts:* 100 grams of pre-prepared *Silybum marianum* seed powder were extracted at room temperature by soaking in 500 ml of petroleum ether for 72 hours, stirring occasionally. The extract was then filtered through Whatman No. 1 filter paper, and the solid residue was extracted using two methods: The sequential method using a variety of solvents (water, ethanol, chloroform, and hexane). The ratio of seed material to extract was (1: 10), and the process was carried out in a Soxhlet apparatus for 10 hours at a temperature of 90°C. The extracted material was then collected in a separate conical flask for each liquid and concentrated under low pressure and temperature using a rotary evaporator [24]. The samples obtained in this way will be used to determine the specific phytochemical composition of the seeds of the *Silybum marianum* L. plant.

Second, add 150 ml of 97.0% ethanol to (40 g) of *Silybum marianum* L. seeds and then heat in a water bath for approximately half an hour. Then cool, filter, and place in a centrifuge. Next, we take the clear liquid from the top of the test tube and place it in a water bath at a temperature of 40 - 50°C. The sample obtained in this way will be used to qualitatively determine the amino acids [25]. Ethanol was deliberately chosen as an extract and solvent due to its polar properties, which enable it to efficiently dissolve a wide variety of active compounds found in plant seeds. Furthermore, it has a positive safety record and environmentally friendly characteristics compared to other polluting and hazardous polar solvents.

*Qualitative tests for detecting the chemically active compounds:* Qualitative phytochemical analysis of extracts derived from the seeds of the plant (*Silybum marianum* L.) was carried out according to the standard procedures and methods described in references [26-29].

*Qualitative tests for amino acids: Ninhydrin reaction:* In a test tube, 1.0 ml of *Silybum marianum* (L.) seed extract was taken, and 1.0 ml of ninhydrin reagent was added. A purple color was observed, indicating the presence of amino acids [25].

*Qualitative determination of amino acids using two-dimensional paper chromatography:* The amino acid content of the sample was estimated using a two-dimensional ascending paper chromatography method in which two liquid systems were used and Whatman No. 1 chromatography paper was used as the stationary phase for the paper chromatography. The first solvent regimen was A -n-butanol: acetic acid: water, at a ratio of 4: 1: 5 (v/v). The second solvent regimen was B- ethanol: toluene, at a ratio of 2: 1 (v/v). Ninhydrin reagent was used as a color developer [25]. The process is carried out in successive steps [30].

*Amino acid separation using one-dimensional paper chromatography:* The amino acids in the seeds of the plant *Silybum marianum* (L.) were separated using a one-dimensional paper chromatography method, using the solvent system: n-butanol: acetic acid: water, in a ratio of 4: 1: 5 (v/v) according to the method mentioned above, and were identified using standard amino acid models [25].

*Separation and identification of amino acids by Thin-Layer Chromatography (TLC):* The amino acids in the seeds of the plant Silybum marianum (L.) were separated by thin-layer chromatography (TLC) using an aluminum-coated silica gel plate (Merk Co. Inc., Kieselgel F256) in a solvent system: n-butanol: acetic acid: water, at a ratio of 4: 1: 5 (v/v), and identified using standard amino acids [25].

*Ethical issues:* No ethical approval is needed for this study.

## Results and discussion

The phytochemicals extracted from the (*Silybum Marianum* L.) plant play an important role in treating many different diseases and are used in the manufacture of many medicinal preparations and nutritional supplements. This plant is still used in traditional and modern folk medicine practices [30]. The results of the phytochemical study conducted on the seeds of *Silybum marianum* L. showed that its seeds contain: flavonoids, amino acids, carbohydrates, sterols, saponins, anthraquinones, phenolic compounds, steroids, terpenoids, tannins, alkaloids, anthocyanins, quinones, and glycosides. The results are shown in **Table 1**.

The current study, which investigated the active compounds in (*Silybum Marianum* L.) seeds, showed good agreement with the results of other studies in Romania, Egypt, Turkey, and Kazakhstan [15, 31-33]. As a result of phytochemical examination (qualitative analysis - chromatographic techniques) and comparison with reference samples, seventeen free amino acids were found in (*Silybum marianum* L.) seeds. Data showed that the seeds of the studied plant contain a wide variety of amino acids: Arginine, leucine, valine, lysine, phenylalanine, isoleucine, threonine, histidine, glutamic acid, glycine, serine, proline, aspartic acid, alanine, tyrosine, methionine, cysteine. The results are shown in **Table 2**. Amino acids, which are found in many parts of plants, are of great importance to plants as they play a vital role in various physiological processes that occur within the plant, such as the synthesis of phenolic compounds and other processes of manufacturing active substances. In addition, those plants that have a large stock of their active components, amino acids, are considered to be an important source of amino acids and can be used in the manufacture of pharmaceutical products, food supplements, and other medical materials [34, 35].

The amino acids lysine and valine are essential for all muscle proteins; tyrosine is necessary for the synthesis of neurotransmitters such as dopamine, norepinephrine, and adrenaline. The amino acid isoleucine is crucial for the production of hemoglobin in red blood cells. Phenylalanine, however, is used in cases of depression and loss of appetite [30]. The results of our current study, which investigated the qualitative chemical analysis of amino acids in *Silybum marianum* L. seeds, showed good agreement with the results of other studies in Uzbekistan, Turkey, Romania, and Poland [21, 32, 36, 37]. The results of the phytochemical analysis of Silybum Marianum L. showed that the seeds contain a variety of active ingredients (**Table 1**), and a large and varied quantity of amino acids (**Table 2**). These results demonstrate the richness of this plant's seeds in active ingredients that play a vital role in bodily functions and possess diverse biological, therapeutic, and pharmacological properties, including antioxidant and antitumor activity, and can be used in the treatment of cancerous diseases [38, 39]. As we mentioned earlier, amino acids play an important and major role in plants through their function as an osmotic factor; they regulate stomatal penetration; they remove toxins, especially heavy metals; and they have a major role in the activity and synthesis of some enzymes [34].

**Table 1:** Qualitative chemical analysis of the active ingredients in the seeds of *Silybum Marianum* L.

| Phytochemicals Compounds | Test | Finding |
|---|---|---|
| **Flavonoids** | Shibata's reaction (Shinoda reaction ) | + |
| **Amino acids** | Biuret and Ninhydrin reaction | + |
| **Carbohydrate** | Molisch's test | + |
| **Sterol** | Liebermann's Burchard | + |
| **Saponin** | Foam (Froth) Test | + |
| **Anthraquinones** | Borntrager's reaction | + |
| **Phenolic compound** | Ferric chloride | + |
| **Steroids and terpenoids** | Salkowski | + |
| **Tannin** | 1.0% Gelatine containing 10% sodium chloride | + |
| **Coumarins** | NaOH solution | - |
| **Alkaloids** | Dragendorff"s Test | + |
| **Anthocyanin** | HCl and ammonia | + |
| **Quinone** | NaOH test | + |
| **Glycosides** | Borntrager's test | + |

**Table 2:** The qualitative composition of amino acids in the seeds of (*Silybum marianum* L.)

| Amino acid | Rf values in mobile phase | |
|---|---|---|
| | A | B |
| **Alanine** | 0.21 | 0.30 |
| **Arginine** | 0.05 | 0.24 |
| **Glutamic acid** | 0.18 | 0.15 |
| **Aspartic acid** | 0.16 | 0.04 |
| **Glycine** | 0.23 | 0.14 |
| **Leucine** | 0.65 | 0.63 |
| **Serine** | 0.18 | 0.11 |
| **Isoleucine** | 0.71 | 0.73 |
| **Methionine** | 0.40 | 0.44 |
| **Threonine** | 0.19 | 0.16 |
| **Cysteine** | 0.46 | 0.03 |
| **Valine** | 0.44 | 0.61 |
| **Lysine** | 0.04 | 0.12 |
| **Histidine** | 0.12 | 0.05 |
| **Proline** | 0.24 | 0.31 |
| **Phenylalanine** | 0.33 | 0.53 |
| **Tyrosine** | 0.55 | 0.37 |

***A-** n-butanol - acetic acid - water (4:1:2)- **B-** ethanol - toluene (2:1)*

The identification and isolation of flavonoid, phenol, and amino acid derivatives in the seeds of the plant (*Silybum Marianum* L.) is linked to the plant's therapeutic properties, and the researcher believes this may be the reason for the use of this plant in traditional Palestinian folk medicine and other countries. However, further studies are still needed on all parts of the plant to identify and isolate the compounds responsible for the plant's distinctive therapeutic properties, especially phenols and flavonoids, which opens the door to the possibility of using its seeds in treating diseases caused by free radicals, such as cancer.

***Conclusions:*** After studying extracts of the *Silybum marianum* L. plant to detect the active ingredients, the results of the study showed that the extracts from the seeds of the plant contain biologically active components such as: flavonoids, amino acids, carbohydrates, sterols, saponin, anthraquinones, phenolic compounds, steroids, terpenoids, tannin, alkaloids, anthocyanin, quinone, and glycosides. For the first time in Palestine, a study conducted on extracts of the seeds of the Palestinian plant (*Silybum Marianum* L.) to detect seventeen amino acids using chemical chromatographic analysis showed that the seeds of this plant contain a variety of essential and non-essential amino acids: arginine, leucine, valine, lysine, phenylalanine, isoleucine, threonine, histidine, glutamic, glycine, serine, proline, aspartic, alanine, tyrosine, methionine, and cysteine.

## References

1. Iraqi O, Aouji M, Oubihi A, Imtara H, Noman O, Mothana R, et al. Exploring the chemical composition, antioxidant, and antibacterial properties of milk thistle (*Silybum marianum* L.). Frontiers in Sustainable Food Systems. 2025; 9: 551584. doi: 10.3389/fsufs.2025.1551584
2. Elshafie HS, Camele I, Mohamed AA. A comprehensive review on the biological, agricultural and pharmaceutical properties of secondary metabolites based on plant origin. International Journal of Molecular Sciences. 2023; 24: 3266. doi: 10.3390/ijms24043266
3. Sidhu MC, Saini P. *Silybum marianum*: A plant of high medicinal importance: A review. World Journal of Pharmaceutical Research. 2012; 1: 72-86. doi: Nil.
4. Marceddu R, Dinolfo L, Carrubba A, Sarno M, Di Miceli G. Milk Thistle (*Silybum marianum* L.) as a novel multipurpose crop for agriculture in marginal environments: A review. Agronomy. 2022; 12(3): 729. doi: 10.3390/agronomy12030729
5. Veres T, Tyr S. Milk thistle (*Silybum marianum* (L.) Gaertn.) as a weed in sustainable crop rotation. Research Journal of Agricultural Sciences. 2012; 44: 118-122. doi: Nil.

6. Ledda L, Deligios P, Farci R, Sulas L. Biomass supply for energetic purposes from some Carduae species grown in a Mediterranean rainfed low input cropping system. Industrial Crops and Products. 2013; 47: 218226. doi: Nil.
7. Ahmad M, Zafar M, Sultana S, Azam A, Khan AM. The optimization of biodiesel production from a novel source of wild non-edible oil yielding plant *Silybum marianum*. International Journal of Green Energy. 2014; 11: 589-594. doi: Nil.
8. Porwal O, Sheet M, Anwer Bayrakdar ET, Uthirapathy S, *Silybum marianum* (milk thistle): Review on its chemistry, morphology, ethno-medical uses, phytochemistry and pharmacological activities. Journal of Drug Delivery and Therapeutics. 2019; 9(5): 199-206. doi: 10.22270/jddt.v9i5.3666
9. Bahmani M, Shirzad H, Rafieian S, Rafieian-Kopaei M. *Silybum marianum*: Beyond hepatoprotection. Journal of Evidence-Based Complementary and Alternative Medicine. 2015; 20(4): 292-301. doi: 10.1177/215658 7215571116
10. Khlaq M, Alum MK, Alam MM. Anti-inflammatory potential of medicinal plants. Mediterranean Journal of Pharmacy and Pharmaceutical Sciences. 2022; 2(1): 13-21. doi: 10.5281/zenodo.6399381
11. Bhat R, Bhat SS, Nisarga P. A systematic review and meta-analysis of traditional Chinese medicine in Parkinsonism. Mediterranean Journal of Medical Research. 2025; 2(4): 169-178. doi: 10.5281/zenodo. 17344302
12. Hoque M, Hasan MN, Saikh S. Using common medicinal plants to treat high blood pressure: An updated overview and emphasis on antihypertensive phytochemicals. Mediterranean Journal of Pharmacy and Pharmaceutical Sciences. 2025; 5(3): 1-10. doi: 10.5281/zenodo.15788473
13. Lee DY-W, Liu Y. Molecular structure and stereochemistry of silybin A, silybin B, isosilybin A, and isosilybin B, isolated from *Silybum marianum* (Milk Thistle). Journal of Natural Products. 2003; 66: 1171.1174. doi: 10.1021/np030163b
14. Deep G, Agarwal R. Antimetastatic efficacy of silibinin: Molecular mechanisms and therapeutic potential against cancer. Cancer Metastasis Reviews. 2010; 29(3): 447-463. doi: 10.1007/s10555-010-9237-0
15. Coșeran MA, Constantinescu E, Ona A, Balaș S, Duda M. Milk Thistle [*Silybum marianum* (L.) Gaertn.] - a very useful plant for people's health. Hop and Medicinal Plants. 2022; 30(1-2): 70-83. doi: 10.15835/hpm.v30i1 -2.14439
16. Smith T, Lynch ME, Johnson J, Kawa K, Bauman H, Blumenthal M. Herbal dietary supplement sales in the US increase 6.8% in 2014. HerbalGram. 2015; 107: 52-59. doi: Nil.
17. Ghaleb T. Quantitative determination of Silymarin in *Silybum marianum*, *Cynara scolymus* and *Gundelia toumefortii* grown in Syria. (A comparative study). Latakia University Journal-Biological Sciences Series. 2025; 46(6): 33-46. doi: Nil.
18. Andrzejewska J, Sadowska K, Mielcarek S. Effect of sowing date and rate on the yield and flavonolignan content of the fruits of milk thistle (*Silybum marianum* L. Gaertn). Industrial Crops and Products. 2011; 33(2): 462-468. doi: 10.1016/j.indcrop.2010.10.027
19. Elwekeel A, Elfishawy A, AbouZid S. Silymarin content in *Silybum marianum* fruits at different maturity stages. Journal of Medicinal Plants Research. 2013; 7: 1665-1669. doi: 10.5897/JMPR12.0743
20. Gurbuz B, Gumuscu A, Arslan N. The effect of plant frequency on seed yield of milk thistle (*Silybum marianum* L.) Proceedings of The XII$^{th}$ Symposium on Plant-Originated Crude Drugs. Ankara, Turkey. 2000; 107-110.
21. Apostol L, Iorga S, Mosoiu C, Mus G, Cucu S. Nutrient composition of partially defatted milk thistle seeds. Scientific Bulletin. Series F. Biotechnologies. 2017; XXI: 165-169. doi: Nil.
22. Abou Auda M. Phytochemical and proximate analysis of wild plants from the Gaza Strip, Palestine. Biodiversitas Journal of Biological Diversity. 2025; 26: 185-194. doi: 10.13057/biodiv/d260307
23. Abou Auda M, Alghuff S, Elbashiti T. Ethnobotanical investigation of medicinal plants and their importance, traded in the public herbal markets and centers of Gaza Strip, Palestine. An-Najah University Journal for Research - A (Natural Sciences). 2023; 27: 21-28. doi: 10.35552/anujr.a.37.1.2059
24. Abu Thaher KTM. The amino acids of leaves and seeds of *Ecballium elaterium* (L.) A. Rich. Flora of Palestine. World Journal of Pharmaceutical and Medical Research. 2024; 10(10): 39-42. doi: Nil.
25. Sankhalkar S, Vernekar V. Quantitative and qualitative analysis of phenolic and flavonoid content in *Moringa oleifera* L. and *Ocimum tenuiflorum* L. Pharmacognosy Research. 2016; 8(1): 16-21. doi: 10.4103/0974-8490. 171095
26. Harborne JB. Phytochemical methods: A guide to modern techniques of plant analysis. .(2008) 3$^{rd}$ ed. London: Chapman and Hall. ISBN: 0 214 57270 2.
27. Kudashkina NV, Khasanova SR, Meshcheryakova SA. Phytochemical analysis: textbook. Allowance (2019) - Ufa: Federal State Budgetary Educational Institution of Higher Education BSMU of the Ministry of Health of Russia. 2019.
28. Mishra T, Kaundal RK. Unveiling the anticancer potential of milk thistle in hepatocellular carcinoma: A network pharmacology perspective. Pharma Nutrition. 2025; 34: 100458. doi: 10.1016/j.phanu.2025.100458

29. Abu Thaher K. Preliminary phytochemical analysis, traditional uses and pharmacological activities of *Solanum nigrum* L. flora of Palestine. International Journal of Pharmacognosy and Pharmaceutical Sciences. 2024; 6: 46-52. doi: 10.33545/27067009.2024.v6.i2a.157
30. Aye MTN, Arnt W, Thi TK, Zin MH. Qualitative and quantitative investigations of amino acids content from the flowers of *Moringa olifera* Lam. (Dan-da-lun). Research Journal. 2020; 11: 2020. University of Mandalay
31. Mahmoud NN, Selim MT. Phytochemical analysis and antimicrobial activity of *Silybum marianum* L. via multi-solvent extraction. AMB Express. 2025; 15(1): 122. doi: 10.1186/s13568-025-01925-2
32. Doğan G, Kara N, Gür S, Bagci E. Chemical composition and biological activity of milk thistle seeds (*Silybum marianum* (L.) Gaertn.). International Journal of Nature and Life Sciences. 2022; 6: 90-98. doi: 10.47947/ijnls.1124
33. Tileshova M, Yessimsiitova Z, Alseitova F, Chunetova Z, Pravin N, Tileubayeva Z, Ryskali T, Yeltay G. Phytochemical study of milk thistle, *Silybum marianum* (L.) Gaertn. Caspian Journal of Environmental Sciences. 2024; 22(5): 1293-1299. doi: 10.22124/cjes.2024.8345
34. Rai V. Role of amino acids in plant responses to stresses. Biologia Plantarum. 2002; 45: 481-487. doi: 10.1023/A:1022308229759
35. Ahmed R, Uddin MM, Hoque M. Nutraceuticals: Food-based therapeutics and health benefits. Mediterranean Journal of Medicine and Medical Sciences. 2025; 1(1): 22-30. doi: 10.5281/zenodo.15771921
36. Kayumov FSU, Oblokulov JB, Jalilov UM, Kayumova GGQ. Amino acid analysis of dry extract of milk thistle fruit *Silybum marianum Gaertn*. L. growing in Uzbekistan. Journal NX. 2021; 7(1): 198-203. doi: Nil.
37. Teleszko M, Haraf G, Zając A, Krzos G. Milk Thistle (*Silybum marianum* (L.) Gaertner) endosperm as an alternative protein source for a sustainable food system (SFS)- Pilot studies. Sustainability. 2023; 15: 14411. doi: 10.3390/su151914411
38. Riaz M, Khalid R, Afzal M, Anjum F, Fatima H, Zia S, et al. Phytobioactive compounds as therapeutic agents for human diseases: A review. Food Science and Nutrition. 2023; 11(6): 2500-2529. doi: 10.1002/fsn3.3308
39. Omer GA, Mohammed LY. Total phenolic, flavonoids and vitamin C contents with antioxidant activity of *Urtica dioica* L. leaves growing in Zakho, Kurdistan Region-Iraq. Baghdad Science Journal. 2024; 21; 5: 18. doi: 10.21123/bsj.2023.8658

**Conflict of interest:** The author declares the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
**Ethical issues:** The author has considered ethical issues, including plagiarism, informed consent, data fabrication or falsification, and double publication or submission.
**Data availability statement:** The raw data that support the findings of this article are available from the corresponding author upon reasonable request.
**Generative AI disclosure:** No Generative AI was used in the preparation of this manuscript.